




ese animals are reared by pastoralist Gujar communities from Haryana and Punjab.

‡ 3` [ƒ[fgf[a` Sl award “Taradevi Modi





Nari Cattle: Nari cattle are found in the Bali

S V _ [j W [^ # ' , S [S ^ ž F Z W

kids are sold for chevon whereas the females are kept as replacements.

For conservation and improvement of livestock genetic resources in the farmers' herds Surti, Gir, Sirohi and Sonadi breeds of buffalo, cattle, goats and sheep species respectively were identified. Seven Gir bulls, 46 Sirohi bucks and 5 Sonadi rams were given to the farmers by NBAGR &

***Ex-situ* conservation**

Genetic Characterization

i [fZ fZWShWdSYMaX"đ %ž3Tagf)' ı aXfZW



5504 position (A>G) in intron- 2 region of the TLR4 gene showed a significant difference in frequencies between mastitis-affected and



Genetic characterization of Red Jungle Fowl

(

Relationships of gene polymorphisms with growth traits in goat: The positive effect of Insulin-like growth factor-1(IGF1), Growth

Goat TLR2 gene has been characterized by polymorphism analysis, which showed presence of seven polymorphic nucleotides in 12 different animals. Out of five SNPs in the coding region, four were synonymous and

Buñalo Genomics

Completed Research Projects

14.

M Mukesh, Rajesh Kumar and AK Tyagi (NDRI)



38.



An Interactive meet

Felicitatation of Kankrej cattle owner on Biodiversity Day



Promotions







fy, vuøe fu/ur fd; k x; k
M[V] l gu'kDr v\$ t\$ jk k fud
ek n. Ma (fØ, fvu d bzt} yDVd



jkt HK'ki izlšB dhxfrfof/k k

fnul 12-7-2011 dksvijkg 3-00 ctsl LFku jkt HK'k dk kZ; u l fefr dh cBd gpZ
ftl eafy, x, egRoivZfu.kZ leal seq; r%C; jks dh oSl kZ dk fgah vuqkn jkt HK'k
izlšB }kjkd; st kusdkfu.kZ gya

fl rEcj 2011 dk 66h 8

